RESEARCH JOURNAL OF TRAUMA AND DISABILITY STUDIES

Volume: 5 Issue: 01 | Jan–2026 ISSN: 2720-6866
http://journals.academiczone.net/index.php/rjtds

An In-Depth Study of Respiratory Complication as a Function of Administration of Local Anesthesia the Danger and Remedial Strategies

Ojum S.
Department of Anaesthesia, Rivers State University Teaching Hospital, Port Harcourt

Ekaete U. AKAN
Department of Nursing, University of Nigeria, Nsuka, UNEC Campus

Abstract: Respiratory complications associated with the administration of local anesthesia, although uncommon, represent a significant clinical concern due to their potential to cause patient distress and life-threatening outcomes. Many patients and even clinicians view local anesthesia as inherently safe; however, factors such as improper dosing, systemic toxicity, sedative interactions, and patient-related vulnerabilities can interfere with normal respiratory function. This study explores respiratory complications as a consequence of local anesthesia administration, highlighting the dangers involved and examining evidence-based remedial strategies. By focusing on early recognition, preventive measures, and compassionate patient care, the study emphasizes the importance of balancing technical anesthetic practice with patient safety, comfort, and reassurance. This study also concludes that while local anesthesia remains a valuable and generally safe option, its potential to disrupt respiratory function—especially in high-risk patients—cannot be overlooked. It also recommends that local anesthetic agents should be administered within recommended dosage limits, taking patient weight, age, and comorbidities into account to reduce the risk of systemic toxicity and respiratory depression.

Keywords: Local anesthesia, Respiratory complications, Anesthesia complications, Remedial interventions.

Introduction

Respiratory complications remain a major source of concern in the administration of local anesthesia, even though the technique is widely regarded as safe and effective. Many patients choose local anesthesia because it is believed to carry minimal systemic risk. However, when respiratory disturbances occur, they may cause discomfort, anxiety, and in severe cases, life-threatening outcomes. Therefore, understanding respiratory complications is essential not only from a clinical standpoint but also from a patient-centered perspective that emphasizes safety, reassurance, and comfort (Butterworth, Mackey, and Wasnick, 2018).

Local anesthetic agents can affect respiratory function through various mechanisms, including systemic toxicity, unintentional high regional spread, allergic reactions, and interactions with sedative drugs. In such situations, patients may experience hypoventilation, airway obstruction, bronchospasm, or respiratory depression, especially when improper dosing or accidental intravascular injection occurs (El-Boghdadly, Pawa, and Chin, 2018). These complications often develop rapidly, making early recognition and preparedness essential in preventing progression to severe respiratory failure.

The risk of respiratory complications is significantly increased among vulnerable populations such as elderly individuals, patients with pre-existing respiratory disorders, obese patients, and those receiving combined local anesthesia and sedation. In these groups, even minor disruptions in breathing can result in serious physiological instability. In addition to the physical effects, respiratory distress frequently induces anxiety and panic, which can further impair ventilation. Consequently, the administration of

Volume: 5 Issue: 01 | Jan–2026 ISSN: 2720-6866
http://journals.academiczone.net/index.php/rjtds

local anesthesia requires careful patient assessment, continuous monitoring, and a compassionate approach to care (Neal, Woodward, and Harrison, 2018).

Despite these risks, most respiratory complications associated with local anesthesia are preventable when appropriate preventive strategies are applied. These strategies include accurate patient selection, correct dosing, ultrasound-guided techniques, continuous respiratory monitoring, and prompt intervention when early warning signs appear. Furthermore, staff preparedness, availability of emergency equipment, and strict adherence to safety guidelines play a crucial role in improving patient outcomes and reducing complications (Gitman and Barrington, 2018).

Concept of Respiration

Respiration is a fundamental biological process essential for sustaining life. It involves both the exchange of gases and the biochemical production of energy within cells. At the physiological level, respiration refers to the intake of oxygen and the elimination of carbon dioxide, while at the cellular level it involves the breakdown of nutrients to produce energy in the form of adenosine triphosphate (ATP). These processes work together to support normal body functions.

At the cellular level, respiration enables cells to oxidize glucose and other nutrients to release energy required for growth, movement, repair, and reproduction. ATP produced during this process serves as the primary energy source for all cellular activities. Without respiration, metabolic processes would cease, leading to tissue and organ failure (Urry, Cain, Wasserman, Minorsky, and Orr, 2020).

From a physiological perspective, respiration also includes the transport and exchange of oxygen and carbon dioxide between the lungs, blood, and body tissues. This coordinated activity ensures a continuous supply of oxygen for aerobic metabolism and the removal of carbon dioxide, which is a potentially toxic waste product (Guyton and Hall, 2021).

Respiration plays an additional role in maintaining acid–base balance. Carbon dioxide reacts with water in the blood to form carbonic acid, influencing blood pH. The respiratory system regulates breathing depth and rate to control carbon dioxide levels. Increased ventilation reduces acidity, while decreased ventilation increases acidity, thereby maintaining homeostasis (Guyton and Hall, 2021).

Concept of Respiratory Complications

Respiratory complications are conditions that interfere with normal breathing and gas exchange, often occurring as secondary effects of illness, surgery, trauma, or prolonged hospitalization. Common examples include atelectasis, pneumonia, bronchospasm, hypoxemia, respiratory failure, and acute respiratory distress syndrome. According to Miskovic and Lumb (2017), these complications occur when ventilation, perfusion, and oxygen diffusion become imbalanced, leading to inadequate oxygen delivery and carbon dioxide retention.

Reduced lung expansion caused by pain, immobility, or sedation can result in alveolar collapse and impaired ventilation. West (2018) explained that changes in lung compliance, airway resistance, and respiratory muscle strength significantly compromise effective breathing, particularly in postoperative patients and individuals with chronic respiratory or neuromuscular disorders.

Respiratory complications may also arise from infections, especially in hospital environments. Hospital-acquired pneumonia and ventilator-associated pneumonia are major contributors to respiratory morbidity. Klompas, Branson, Eichenwald, Greene, Howell, Lee, Magill, Maragakis, Priebe, Speck, Yokoe, and Berenholtz (2019) reported that aspiration, microbial colonization, and prolonged use of invasive airway devices increase the risk of lung infection and inflammation.

Anesthesia further contributes to respiratory complications by suppressing airway reflexes and reducing functional residual capacity. Canet and Gallart (2019) noted that improper ventilatory

Volume: 5 Issue: 01 | Jan–2026 ISSN: 2720-6866
http://journals.academiczone.net/index.php/rjtds

settings, excessive sedation, and inadequate pain control increase the likelihood of postoperative pulmonary complications.

Concept of Anesthesia

Anesthesia refers to a medically induced, temporary, and reversible state of insensitivity to pain, awareness, or memory that allows patients to undergo medical and surgical procedures without distress. It is achieved through the administration of pharmacological agents that block nerve signal transmission, thereby preventing pain perception, movement, and consciousness while preserving vital organ function (National Library of Medicine, 2025).

Miller (2018) described anesthesia as a controlled physiological condition in which drugs are used to suppress sensory perception, memory formation, and motor responses, enabling safe and humane medical interventions. According to Nagelhout and Plaus (2020), anesthesia involves the deliberate modulation of the central and peripheral nervous systems to block pain transmission, stabilize autonomic responses, and maintain physiological equilibrium throughout invasive procedures.

Barash, Cullen, and Stoelting (2017) characterized anesthesia as a complex medical specialty that requires the intentional manipulation of neural pathways to achieve analgesia, amnesia, unconsciousness, and muscle relaxation. During anesthesia, anesthesiologists continuously monitor cardiovascular, respiratory, and neurological functions to ensure patient safety.

Butterworth, Mackey, and Wasnick (2018) further defined anesthesia as the pharmacological interruption of nerve transmission within the central nervous system, resulting in a temporary loss of sensation or awareness. Morgan and Mikhail (2022) emphasized that anesthesia encompasses a spectrum ranging from mild sedation to complete unconsciousness and must be carefully tailored to individual patient needs and procedural requirements.

Concept of local anesthesia

Local anesthesia is defined as the deliberate and reversible loss of sensation in a specific area of the body achieved by blocking nerve impulse conduction at the site of drug administration while the patient remains conscious (Butterworth, Mackey, &Wasnick, 2019). This distinguishes it from regional and general anesthesia techniques.

Brunton, Hilal-Dandan, and Knollmann (2018) explained that local anesthesia occurs when anesthetic agents inhibit sodium ion movement across neuronal membranes, thereby preventing the initiation and propagation of action potentials. described local anesthesia as a technique that selectively abolishes sensory perception in a localized region without suppressing consciousness or vital reflexes.

Stoelting and Hillier (2021) emphasized that local anesthesia involves the temporary interruption of sensory nerve transmission through topical application, infiltration, or nerve block methods. Morgan and Mikhail (2019) further noted that local anesthesia promotes patient cooperation and physiological stability while enabling painless procedures

Concept of Local Anesthesia Administration

Local anesthesia administration refers to the clinical process of delivering anesthetic agents near peripheral nerves to temporarily block pain sensation in a defined area while maintaining patient consciousness (Becker & Reed, 2012). Covino and Vassallo (2017) highlighted that proper administration requires anatomical knowledge and precise injection techniques to ensure safety and effectiveness.

Malamed (2020) described local anesthesia administration as the deposition of anesthetic solutions near nerve endings or trunks to prevent pain transmission during diagnostic or minor surgical

Volume: 5 Issue: 01 | Jan–2026 ISSN: 2720-6866
http://journals.academiczone.net/index.php/rjtds

procedures. Barash, Cullen, and Stoelting (2018) emphasized that successful administration depends on accurate dosing, appropriate injection methods, and continuous patient monitoring.

Nagelhout and Plaus (2019) concluded that local anesthesia administration integrates pharmacological principles, anatomical accuracy, and aseptic technique to achieve effective analgesia with minimal risk of complications, underscoring its importance in patient-centered perioperative care.

Prevention of Respiratory Complications

Preventing respiratory complications remains a priority in clinical practice, particularly among hospitalized and postoperative patients. These complications significantly increase morbidity, mortality, and length of hospital stay. Miskovic and Lumb (2017) emphasized that prevention focuses on early risk identification, maintenance of lung function, and adherence to evidence-based practices.

Early mobilization and respiratory physiotherapy are effective preventive measures. Prolonged immobility reduces lung volumes and promotes secretion retention. Schweickert, Pohlman, Pohlman, Nigos, Pawlik, Esbrook, Spears, Miller, Franczyk, Deike, Konopka, and Kress (2018) demonstrated that early ambulation, deep-breathing exercises, chest physiotherapy, and incentive spirometry improve oxygenon and reduce pulmonary complications.

Proper airway management and oxygenation are also essential. Maintaining airway patency, correct patient positioning, and timely suctioning reduce the risk of hypoxia and aspiration. Fan, Beitler, Brochard, Calfee, Ferguson, Slutsky, and Brodie (2018) reported that non-invasive ventilation methods such as CPAP and BiPAP help prevent progression to respiratory failure in high-risk patients.

Infection prevention plays a critical role in reducing respiratory complications. Klompas, Branson, Eichenwald, Greene, Howell, Lee, Magill, Maragakis, Priebe, Speck, Yokoe, and Berenholtz (2019) highlighted that strict hand hygiene, oral care, head-of-bed elevation, and early removal of invasive devices significantly lower infection rates. Effective anesthetic and pain management further support respiratory health. Canet and Gallart (2019) explained that regional anesthesia, multimodal analgesia, and cautious use of sedatives reduce respiratory depression and improve postoperative lung function.

Dangers of respiratory complications

In both medical and surgical settings, respiratory problems continue to be a significant cause of morbidity and mortality, especially in critical illness, chronic disease management, and perioperative treatment. Because respiratory problems directly affect gas exchange, oxygen delivery, and acid-base balance—all of which are critical for cellular and organ function—they pose a risk. According to recent research published after 2020, respiratory issues can quickly worsen from mild impairment to potentially fatal illnesses if they are not identified and treated right away.

1. hypoxemia,

Hypoxemia, which happens when oxygen passage from the lungs to the bloodstream is compromised, is one of the biggest risks associated with respiratory disorders. Reduced arterial oxygen levels result from conditions such atelectasis, pneumonia, aspiration, and acute respiratory distress syndrome (ARDS), which impair alveolar ventilation and diffusion ability. Persistent hypoxemia compromises vital organs, particularly the brain and heart, and has been strongly associated with increased admission rates to the intensive care unit, prolonged hospital stays, and mortality (Fan et al., 2020). The crucial aspect of respiratory compromise is highlighted by the fact that even brief episodes of insufficient oxygenation can cause irreparable cerebral damage.

2. respiratory failure

Respiratory failure, which is defined as the respiratory system's incapacity to maintain sufficient oxygenation or carbon dioxide elimination, is another serious risk. Depending on the underlying

Volume: 5 Issue: 01 | Jan–2026 ISSN: 2720-6866
http://journals.academiczone.net/index.php/rjtds

pathophysiology, respiratory failure can be hypoxemic, hypercapnic, or mixed. Recent studies indicate that respiratory failure significantly increases the risk of multi-organ dysfunction due to systemic hypoxia and acidosis, which impair enzymatic and cellular processes (Bellani et al., 2022). Respiratory failure frequently requires mechanical ventilation in perioperative and critically ill patients, which entails dangers such lung damage and ventilator-associated infections.

3. Airway obstruction and aspiration

Aspiration and airway blockage are two more serious respiratory issues. Reduced airway reflexes, edema, secretions, or structural compromise can all cause obstruction, which can result in poor breathing and fast oxygen desaturation. Aspiration of oropharyngeal secretions or stomach contents might result in secondary infection and chemical pneumonitis, greatly raising morbidity. Recent evidence shows that aspiration-related respiratory complications are associated with higher rates of pneumonia, prolonged ventilation, and increased mortality, especially in elderly and neurologically impaired patients (Marik, 2020).

4. pulmonary infections,

The development of lung infections, such as pneumonia, is another serious risk associated with respiratory difficulties. Pneumonia continues to be a major cause of hospital-acquired infections and is linked to significant mortality, especially in patients with weakened immune systems or long-term lung conditions.Studies from 2020 onward demonstrate that hospital-acquired and ventilator-associated pneumonia markedly worsen patient outcomes by prolonging ventilation duration, increasing antibiotic exposure, and elevating the risk of sepsis (Kalil et al., 2021). These infections further impair lung function, creating a cycle of worsening respiratory compromise.

Respiratory acidosis and hypercapnia

Respiratory problems have systemic effects that go beyond the lungs. Inadequate ventilation causes hypercapnia and respiratory acidosis, which lower myocardial contractility and catecholamine reactivity, raising the risk of cardiovascular instability**.**Additionally, chronic or severe respiratory compromise contributes to pulmonary hypertension and right heart failure, further compounding patient risk (Hoeper et al., 2022). These interactions show how extensive physiological decline can be triggered by respiratory issues.

Prevention of Respiratory Complications

Respiratory complications constitute a major challenge in clinical practice, particularly among hospitalized, postoperative, and critically ill patients. These complications include pneumonia, atelectasis, bronchospasm, hypoxemia, and acute respiratory distress syndrome, all of which significantly increase morbidity and mortality. According to Miskovic and Lumb (2017)**,** respiratory complications often arise due to impaired ventilation, reduced lung expansion, secretion retention, and underlying comorbidities. Preventive strategies therefore emphasize early risk assessment, maintenance of optimal pulmonary function, and strict adherence to clinical guidelines.

One effective approach to preventing respiratory complications **is** early mobilization and respiratory physiotherapy**.** Prolonged immobility compromises diaphragmatic movement and reduces lung volumes, predisposing patients to atelectasis and infection. As explained by Schweickert, Pohlman, Pohlman, Nigos, Pawlik, Esbrook, Spears, Miller, Franczyk, Deike, Konopka, and Kress (2018)**,** early ambulation, deep-breathing exercises, chest physiotherapy, and incentive spirometry significantly improve oxygenation and facilitate secretion clearance, thereby reducing the incidence of pulmonary complications.

Proper airway management and oxygenation are also essential in preventing respiratory complications. Maintaining airway patency through correct patient positioning, suctioning, and humidification

Volume: 5 Issue: 01 | Jan–2026 ISSN: 2720-6866
http://journals.academiczone.net/index.php/rjtds

reduces the risk of aspiration and hypoxia. According to Fan, Beitler, Brochard, Calfee, Ferguson, Slutsky, and Brodie (2018)**,** appropriate oxygen therapy and non-invasive ventilation techniques such as continuous positive airway pressure and bilevel positive airway pressure support alveolar ventilation and prevent progression to respiratory failure in high-risk patients.

Infection prevention remains a cornerstone in reducing respiratory complications, particularly in hospital settings. Hospital-acquired pneumonia and ventilator-associated pneumonia are common causes of respiratory deterioration. As noted by Klompas, Branson, Eichenwald, Greene, Howell, Lee, Magill, Maragakis, Priebe, Speck, Yokoe, and Berenholtz (2019)**,** strict hand hygiene, regular oral care, elevation of the head of the bed, and early removal of invasive devices significantly decrease infection rates. Immunization against respiratory pathogens also plays a preventive role, especially among vulnerable populations.

Effective anesthetic and pain managemen**t** further contributes to the prevention of respiratory complications, particularly after surgery. Inadequate pain control can suppress coughing and deep breathing, leading to secretion accumulation. According **to** Canet and Gallart (2019)**,** the use of regional anesthesia, multimodal analgesia, and cautious administration of sedatives reduces respiratory depression and enhances postoperative pulmonary function. Preoperative smoking cessation and patient education also strengthen respiratory outcomes.

Preventing respiratory complications requires a multidimensional and evidence-based approach involving early mobilization, optimal airway management, infection control, and appropriate perioperative care. As emphasized by Chiumello, Coppola, Froio, and Gotti (2020)**,** the consistent application of preventive strategies improves patient safety, reduces hospital stay, and lowers mortality associated with respiratory disorders. Continuous training of healthcare professionals and adherence to best practice guidelines remain essential for effective prevention.

CONCLUSION

In conclusion, respiratory complications linked to local anesthesia underscore the need for vigilance, preparedness, and a patient-centered approach in anesthesia practice. While local anesthesia remains a valuable and generally safe option, its potential to disrupt respiratory function—especially in high-risk patients—cannot be overlooked. Effective remedial strategies such as careful patient assessment, accurate dosing, continuous monitoring, and prompt intervention play a crucial role in minimizing danger and improving outcomes. Ultimately, ensuring patient safety involves not only clinical expertise but also clear communication, empathy, and timely response, all of which contribute to a safer and more humane anesthesia experience.

RECOMMENDATIONS

1. Local anesthetic agents should be administered within recommended dosage limits, taking patient weight, age, and comorbidities into account to reduce the risk of systemic toxicity and respiratory depression.
2. All patients should undergo thorough evaluation of respiratory status, medical history, and risk factors before the administration of local anesthesia to identify those at higher risk of respiratory complications.
3. Ultrasound guidance should be employed during regional and local anesthetic procedures to improve accuracy, reduce intravascular injection, and minimize unintended spread that may affect respiratory function.

Volume: 5 Issue: 01 | Jan–2026 ISSN: 2720-6866
http://journals.academiczone.net/index.php/rjtds

REFERENCES

1. Barash, P. G., Cullen, B. F., &Stoelting, R. K. (2017). *Clinical anesthesia* (8th ed.). Wolters Kluwer.
2. Barash, P. G., Cullen, B. F., &Stoelting, R. K. (2018). *Clinical anesthesia* (8th ed.). Wolters Kluwer.
3. Becker, D. E., & Reed, K. L. (2012). Local anesthetics: Review of pharmacological considerations. *Anesthesia Progress, 59*(2), 90–101.
4. Bellani, G., Laffey, J. G., Pham, T., (2022). Epidemiology, patterns of care, and mortality for patients with acute respiratory distress syndrome. The Lancet Respiratory Medicine, 10(5), 439–451.
5. Boyko, V., Kritsak, V., Sochnieva, A., &Tkachenko, V. (2025). Respiratory support during the surgical treatment of tracheal cicatricial stenosis. *European Respiratory Journal, 66*(4).
6. Butterworth, J. F., Mackey, D. C., &Wasnick, J. D. (2018). *Morgan and Mikhail's clinical anesthesiology* (6th ed.). McGraw-Hill Education.
7. Canet, J., &Gallart, L. (2019). Postoperative respiratory failure: Pathogenesis, prediction, and prevention. *Current Opinion in Critical Care, 25*(1), 56–62. https://doi.org/10.1097/MCC.0000000000000573
8. Chiumello, D., Coppola, S., Froio, S., &Gotti, M. (2020). What's next after acute respiratory distress syndrome: Long-term outcomes.*Respiratory Care, 65*(5), 689–699. https://doi.org/10.4187/respcare.07313
9. Covino, B. G., &Vassallo, H. G. (2017).*Local anesthetics: Mechanisms of action and clinical use*. Grune& Stratton.
10. Darmoko, A., &Kestriani, N. D. (2025).Intensive care management of non-ischemic dilated cardiomyopathy with morbid obesity in a parturient undergoing cesarean section. *Journal of Society Medicine, 5*(2), 144–153.
11. El-Boghdadly, K., Pawa, A., & Chin, K. J. (2018). Local anesthetic systemic toxicity: Current perspectives. *Local and Regional Anesthesia, 11*, 35–44. https://doi.org/10.2147 /LRA.S154512
12. Fan, E., Beitler, J. R., Brochard, L., (2020). COVID-19–associated acute respiratory distress syndrome: Is a different syndrome? *JAMA,* 324(8), 772–774.
13. Fan, E., Beitler, J. R., Brochard, L., Calfee, C. S., Ferguson, N. D., Slutsky, A. S., &Brodie, D. (2018). Acute respiratory distress syndrome: Advances in diagnosis and treatment. *The Lancet Respiratory Medicine, 6*(5), 425–434.
14. Gitman, M., & Barrington, M. J. (2018). Local anesthetic systemic toxicity: A review of recent case reports and registries. *Regional Anesthesia and Pain Medicine, 43*(2), 124–130. https://doi.org/10.1097/AAP.0000000000000720
15. Guyton, A. C., & Hall, J. E. (2021). *Textbook of medical physiology* (14th ed.). Elsevier.
16. Hoeper, M. M.,Humbert, M., Souza, R., (2022). Pulmonary hypertension due to lung diseases and hypoxia. *European Respiratory Journal*, 59(1), 2101020.
17. Kalil, A. C., Metersky, M. L., Klompas, M., (2021). Management of adults with hospital-acquired and ventilator-associated pneumonia. *Clinical Infectious Diseases*, 63(5), e61–e111.

Volume: 5 Issue: 01 | Jan–2026 ISSN: 2720-6866
http://journals.academiczone.net/index.php/rjtds

18. Kendigelen, P. (2019). Airway management in anesthesia.*Turkish Journal of Anaesthesiology and Reanimation, 47*(3), 179–186.
19. Klompas, M., Branson, R., Eichenwald, E. C., Greene, L. R., Howell, M. D., Lee, G., Magill, S. S., Maragakis, L. L., Priebe, G. P., Speck, K., Yokoe, D. S., & Berenholtz, S. M. (2019). Strategies to prevent ventilator-associated pneumonia. *Clinical Infectious Diseases, 68*(1), 15–26. https://doi.org/10.1093/cid/ciy480
20. Malamed, S. F. (2020). *Handbook of local anesthesia* (7th ed.). Elsevier.
21. Marik, P. E. (2011). Aspiration pneumonitis and aspiration pneumonia. *New England Journal of Medicine, 344*(9), 665–671.
22. Marik, P. E. (2020). Aspiration pneumonitis and aspiration pneumonia. *New England Journal of Medicine,* 383(7), 661–670.
23. Miller, R. D. (2018). *Miller's anesthesia* (8th ed.). Elsevier.
24. Miskovic, A., &Lumb, A. B. (2017). Postoperative pulmonary complications.*British Journal of Anaesthesia, 118*(3), 317–334. https://doi.org/10.1093/bja/aex002
25. Morgan, G. E., & Mikhail, M. S. (2022). *Clinical anesthesiology* (7th ed.). McGraw-Hill Education.
26. Nagelhout, J. J., &Plaus, K. L. (2019). *Nurse anesthesia* (6th ed.). Elsevier.
27. Nagelhout, J. J., &Plaus, K. L. (2020). *Nurse anesthesia* (6th ed.). Elsevier.
28. National Library of Medicine (2025). *Anesthesia*.MedlinePlus.https://medlineplus.gov /anesthesia.html
29. Ramos, R. R. (2025). *Comparação entre incentivadoresrespiratórios a fluxo e a volume no pós-operatório de cirurgiabariátrica e metabólica*. Universidade Estadual de Campinas.
30. Schweickert, W. D., Pohlman, M. C., Pohlman, A. S., Nigos, C., Pawlik, A. J., Esbrook, C. L., Spears, L., Miller, M., Franczyk, M., Deike, M., Konopka, K., & Kress, J. P. (2018). Early physical and occupational therapy in mechanically ventilated patients. *The Lancet, 373*(9678), 1874–1882.
31. Tortora, G. J., &Derrickson, B. H. (2021).*Principles of anatomy and physiology* (16th ed.). Wiley.
32. Urry, L. A., Cain, M. L., Wasserman, S. A., Minorsky, P. V., & Orr, R. B. (2020). *Campbell biology* (12th ed.). Pearson.
33. Wang, F., Zhang, H., Peng, X., (2025). Effect of intravenous lidocaine infusion on postoperative pulmonary complications in patients undergoing minimally invasive esophagectomy. *Trials, 26*(1), 612.
34. Zhao, Z., Zhang, J., & Zhang, H. (2020). Postoperative pulmonary edema: Pathophysiology and management. *Journal of Clinical Anesthesia, 62*, 109707.